

Definable proper actions and equivariant definable Tietze extension

Tomohiro Kawakami

Department of Mathematics, Faculty of Education, Wakayama University,
Sakaedani Wakayama 640-8510, Japan

Abstract

Let $\mathcal{N} = (R, +, \cdot, <, \dots)$ be an o-minimal expansion of the standard structure of a real closed field R . Let G be a definable group and X a definable proper definable G set. We prove that X has only finitely many orbit types. We also prove equivariant definable Tietze extension theorem.

2010 *Mathematics Subject Classification*. 14P10, 57S10, 03C64.

Keywords and Phrases. Finiteness of orbit types, equivariant definable Tietze extension theorem, o-minimal, real closed fields.

1. Introduction.

Let $\mathcal{N} = (R, +, \cdot, <, \dots)$ be an o-minimal expansion of the standard structure of a real closed field R .

General references on o-minimal structures are [2], [3], and also see [9]. Any definable category is a generalization of the semialgebraic category and the definable category on $\mathcal{R} = (R, +, \cdot, <)$ coincides with the semialgebraic one. It is known in [8] that there exist uncountably many o-minimal expansions of the field \mathbb{R} of real numbers.

Let G be a definable group. A *definable G set* means a pair consisting of a definable set X and a group action $\phi : G \times X \rightarrow X$ such that ϕ is definable. A definable map between definable sets is called *definably proper* if the inverse image of every definably compact definable set is definably compact. We call a definable G set X a *proper definable G set* if the map $G \times X \rightarrow X \times$

X defined by $(g, x) \mapsto (gx, x)$ is definably proper.

Let G be a definable group. We can define *orbit types* as well as G is definably compact ([5]).

Theorem 1.1. *Let G be a definable group. Then every proper definable G set X has only finitely many orbit types.*

Theorem 1.1 is proved the case where R is the field \mathbb{R} of real numbers ([5]).

The following theorem is an equivariant version of definable Tietze extension theorem [1]

Theorem 1.2. *Let G be a definably compact definable group, X a definable G set and A a G invariant definably compact definable subset of X . Every G invariant definable function $f : A \rightarrow R$ is extensible to a G invariant definable function $F : X \rightarrow R$ with $F|_A = f$.*

2. Proof of results.

Let $X \subset R^n$ and $Y \subset R^m$ be definable sets. A continuous map $f : X \rightarrow Y$ is *definable* if the graph of f ($\subset X \times Y \subset R^n \times R^m$) is a definable set. A group G is a *definable group* if G is a definable set and the group operations $G \times G \rightarrow G$ and $G \rightarrow G$ are definable. A definable subset X of R^n is *definably compact* if for every definable map $f : (a, b)_R \rightarrow X$, there exist the limits $\lim_{x \rightarrow a+0} f(x)$, $\lim_{x \rightarrow b-0} f(x)$ in X , where $(a, b)_R = \{x \in R \mid a \leq x < b\}$, $-\infty \leq a < b \leq \infty$. A definable subset X of R^n is *definably compact* if and only if X is closed and bounded ([7]). Note that if X is a definably compact definable set and $f : X \rightarrow Y$ is a definable map, then $f(X)$ is definably compact.

We say that two homogeneous proper definable G sets are *equivalent* if they are definably G homeomorphic. Let (G/H) denote the equivalence class of G/H . The set of all equivalence classes of homogeneous proper definable G sets has a natural order defined as $(X) \geq (Y)$ if there exists a definable G map $X \rightarrow Y$. By the definition the reflexivity and the transitivity clearly hold. If $(X) = (G/H)$ and $(Y) = (G/K)$, then $(X) \geq (Y)$ if and only if H is conjugate to a definable subgroup of K . By a way similar to the proof of 4.1 [5], we have the following lemma.

Lemma 2.1. *Let G be a definable group, H a definable subgroup of G and $g \in G$. If $gHg^{-1} \subset H$, then $gHg^{-1} = H$.*

By Lemma 2.1, the anti-symmetry is true.

By a way to similar to the proof of 1.1 [5], we have Theorem 1.1.

Note that every definable subgroup of a definable group is closed ([6]) and a closed subgroup of a definable group is not necessarily definable. For example \mathbb{Z} is a closed subgroup of \mathbb{R} but not a definable subgroup of \mathbb{R} .

Recall existence of definable quotient.

Theorem 2.2 (Existence of definable quotient, 10.2.18 [2]). *Let G be a definably compact definable group and X a*

definable G set. Then the orbit space X/G exists as a definable set, and the orbit map $\pi : X \rightarrow X/G$ is definable, surjective and definably proper.

The following theorem is the topological case of Tietze extension theorem.

Theorem 2.3 (Tietze extension theorem). *Let X be a normal space and A a closed subset of X . Then every continuous map $f : A \rightarrow \mathbb{R}$ is extensible to a continuous map $F : X \rightarrow \mathbb{R}$ with $F|_A = f$.*

The following theorem is the definable case of Tietze extension theorem.

Theorem 2.4 (Definable Tietze extension theorem, [1]). *Let A be a definable closed subset of R^n . Then every definable map $f : A \rightarrow R$ is extensible to a definable map $F : R^n \rightarrow R$ with $F|_A = f$.*

A definable map $f : X \rightarrow Y$ is *definably closed* if for any definable closed subset A of X , $f(A)$ is a definable closed subset of Y .

Theorem 2.5 ([4]). *Let $f : X \rightarrow Y$ be a definable map. Then f is definably proper if and only if f is definably closed and has definably compact fibers.*

Proof of Theorem 1.2. By Theorem 2.1, X/G exists as a definable set in R^n and the projection $\pi : X \rightarrow X/G$ is a surjective definable definably proper map. By Theorem 2.4 and A is definably compact, $\pi(A)$ is a definable closed subset of R^n . Since f is a G invariant definable map, it induces a definable map $f' : f(A) \rightarrow R$ with $f = \pi \circ f'$. By Theorem 2.2, there exists a definable map $F : R^n \rightarrow R$ with $F|_{f(A)} = f'$. Hence $H = \pi \circ F$ is the required map. \square

References

- [1] M. Aschenbrenner and A. Fischer, *Definable versions of theorems by Kirszbraun and Helly*, Proc. Lond. Math. Soc. **102** (2011), 468–502.

- [2] L. van den Dries, *Tame topology and o-minimal structures*, Lecture notes series **248**, London Math. Soc. Cambridge Univ. Press (1998).
- [3] L. van den Dries and C. Miller, *Geometric categories and o-minimal structures*, Duke Math. J. **84** (1996), 497-540.
- [4] M. Edmundo, M. Mamino and L. Prelli, *On definably proper maps*, arXiv:1404.6634.
- [5] T. Kawakami, *Definable C^r groups and proper definable actions*, Bull. Fac. Ed. Wakayama Univ. Natur. Sci. **58** (2008), 9–18.
- [6] A. Pillay, *On groups and fields definable in o-minimal structures*, J. Pure Appl. Algebra **53** (1988), 239-255.
- [7] Y. Peterzil and C. Steinhorn, *Definable compactness and definable subgroups of o-minimal groups*, J. London Math. Soc. **59** (1999), 769–786.
- [8] J.P. Rolin, P. Speissegger and A.J. Wilkie, *Quasianalytic Denjoy-Carleman classes and o-minimality*, J. Amer. Math. Soc. **16** (2003), 751-777.
- [9] M. Shiota, *Geometry of subanalytic and semialgebraic sets*, Progress in Math. **150** (1997), Birkhäuser.